

# Multidisciplinary approach to assessment and intervention of feeding problems in children with autism spectrum disorders: a clinical perspective

*Isabelle Beaudry-Bellefeuille*

Clinica de Terapia Ocupacional Pediátrica Beaudry-Bellefeuille, Oviedo, Spain

*Maria Pomoni*

School of Health Sciences, Faculty of Education, Health and Human Sciences, University of Greenwich, London, UK

*Angharad Welch*

Independent SLT, Find the Key Speech and Language Therapy, Herefordshire, UK

*Tania Moriyón-Iglesias*

Clinica de Terapia Ocupacional Pediátrica Beaudry-Bellefeuille, Oviedo, Spain

*Marta Suárez-González*

Hospital Universitario Central de Asturias, Oviedo, Spain, and

*Eduardo Ramos-Polo*

Private Practice, Oviedo, Spain

## Abstract

**Purpose** – The aim of this paper is to share the details of a multidisciplinary approach, which includes occupational therapy, and to review the factors that should be considered in the evaluation and treatment of children with autism spectrum disorders (ASD) who are excessively selective in their food choices. Issues in this area are complex and often related to several complementary domains (medical, nutritional, psychosocial, sensorimotor, etc.). However, feeding disorders are frequently assessed and treated from a single discipline and important issues are missed or confounded.

**Design/methodology/approach** – A team of experienced clinicians in the field of paediatric feeding disorders gathered the knowledge and experience they acquired from working with individuals with ASD as well as with individuals with other neurodevelopmental diagnosis. A review of current literature in paediatric feeding disorders was used to document and explicate the multifactorial nature of feeding disorders in children with ASD and justify the need for a multidisciplinary approach to issues in this area.

**Findings** – Feeding disorders in children with ASD are linked to multiple sensory, motor, behavioural, nutritional and gastrointestinal comorbidities. A multidisciplinary approach is needed and increasingly recommended. However, multidisciplinary teams, specialised in the care of children with ASD and feeding issues, continue to be difficult to locate and access for families. The authors sought to highlight the signs of feeding problems in children with ASD from different domains and share a model of a multidisciplinary approach that can lead to more successful interventions.

**Originality/value** – The detailed description of the domains linked to feeding issues and the clinical descriptions provided throughout the paper create a roadmap for other clinicians aiming to set up similar teams.

**Keywords** Occupational therapy, Multidisciplinary, Autism, Feeding

**Paper type** Technical paper

## Background

Autism spectrum disorders (ASD) are neurodevelopmental disorders that are characterised by persistent challenges in social interaction, communication and restricted/repetitive

The current issue and full text archive of this journal is available on Emerald Insight at: <https://www.emerald.com/insight/2398-8819.htm>



Irish Journal of Occupational Therapy  
49/2 (2021) 77–83  
Emerald Publishing Limited [ISSN 2398-8819]  
[DOI 10.1108/IJOT-12-2020-0019]

© Isabelle Beaudry-Bellefeuille, Maria Pomoni, Angharad Welch, Tania Moriyón-Iglesias, Marta Suárez-González and Eduardo Ramos-Polo. Published in *Irish Journal of Occupational Therapy*. Published by Emerald Publishing Limited. This article is published under the Creative Commons Attribution (CC BY 4.0) licence. Anyone may reproduce, distribute, translate and create derivative works of this article (for both commercial and non-commercial purposes), subject to full attribution to the original publication and authors. The full terms of this licence may be seen at <http://creativecommons.org/licenses/by/4.0/legalcode>

*Author Declaration and Conflict of interest:* Authors have no conflicts of interest to declare.

*Funding and Acknowledgement:* No funding was received for this paper.

Received 8 December 2020

Revised 8 March 2021

10 April 2021

22 June 2021

21 July 2021

Accepted 9 August 2021

behaviours (American Psychiatric Association, 2013). Feeding problems have been observed in children with ASD from the earliest diagnostic description (Kanner, 1943) and continue to be reported in many children with ASD (Sharp et al., 2013). During the past two decades, feeding problems have been identified as a common co-existing set of problems in individuals with ASD (Sharp et al., 2013). Atypical behaviours related to feeding tend to have an early onset in children with ASD, often being reported within the first year of life (Emond et al., 2010).

One of the most reported reasons for referral of children with ASD to feeding clinics is extreme food selectivity (Bandini et al., 2010), defined as the intake of a limited variety of foods. Children with ASD can follow a restricted diet consisting of five to six or fewer food items or refuse all foods from one or more food groups (Sharp et al., 2018). Selectivity can be based on the texture, colour, taste or smell of food as well as on strong preferences for certain commercial brands (Mari-Bauset et al., 2014). Although these aberrant eating behaviours are common in typically developing children, in children with ASD, they appear to be more frequent, take more time to overcome or even persist into adulthood (Kuschner et al., 2015). In some cases, children with ASD present with obsessive behaviours such as wanting to drink only from a certain cup or eat only from a certain plate (Nadon et al., 2011).

Restricted diet in the ASD population can lead to health problems such as being underweight (Mari-Bauset et al., 2015), overweight (Broder-Fingert et al., 2014) or feeling too tired to effectively attend the school (Florence et al., 2008). Nadon et al. (2011) found that children with ASD were more likely to skip eating at day care, school, family outings to restaurants or with extended family and friends than their siblings, missing out on the multiple benefits of eating in the company of significant others.

The prevalence of eating problems in ASD is reported to be as high as 90% (Kodak and Piazza, 2008). Twachtman-Reilly et al. (2008) reported that 70% of the ASD paediatric population could be described as selective eaters. Given the complex nature of eating problems in children with ASD, a heterogeneous patient population known to have multiple sensory, behavioural, nutritional and gastrointestinal (GI) comorbidities, a multidisciplinary approach is needed and increasingly recommended (Smile et al., 2020). However, multidisciplinary teams specialising in the care of children with ASD and in eating problems are difficult to locate and access for families (Smile et al., 2020).

The aim of this paper is to share the details of a multidisciplinary approach, which includes occupational therapy, and that has proven to be effective in the assessment and treatment of children with ASD and other neurodevelopmental diagnosis who refuse to eat or who are excessively selective in their food choices (Beaudry-Bellefeuille et al., 2015; Gándara-Gafo et al., 2021). The factors to be considered in the evaluation and intervention of selective eating are presented; the roles of the team members are discussed; a model of multidisciplinary collaboration is proposed.

Our multidisciplinary approach is essentially a network approach in which each professional works from their own location (office, clinic, etc.), and all members possess basic knowledge of the areas of expertise of their colleagues.

As experienced clinicians in the field of paediatric feeding difficulties (one gastroenterologist, two occupational therapists [OT], one nutritionist, one speech therapist and 1 psychology researcher specialized in eating problems in children with ASD), we gathered the knowledge and experience acquired from working with individuals with ASD as well as with individuals with other neurodevelopmental disorders. Through discussions and a qualitative review of pertinent literature from each professional domain, the multifactorial nature of eating problems in children with ASD was documented and explicated to support the multidisciplinary approach used by our team.

### Factors to consider when working with children with autism spectrum disorders with feeding issues


Discussions within our team and review of the literature led to a selection of factors collectively considered within our clinical practices. Factors were included when any member of the team considered them important to their area of practice and could provide supportive literature.

#### Motive for consultation

Identifying and analysing parental concerns is the first stage of the assessment process. Any concern related to feeding should be taken seriously, and parents may need to be referred for professional advice (Kerzner et al., 2015). Issues may range anywhere from the family needing basic information on the normal development of feeding to serious medical, nutritional, psychosocial or feeding skills problems that need to be specifically addressed (Kerzner et al., 2015).

Regardless of the initial concern of the family or the first professional to assess the child, a fluid and non-hierarchical relationship and referral system between team members, with the child and the family at the forefront of the process, have been identified as key to successful interventions by our team (Figure 1). For example, a family may initially consult with the

**Figure 1** A fluid and non-hierarchical relationship between the members of the team has been identified as key to successful interventions



psychologist, concerned with their child's behaviour at mealtimes. However, once the psychologist has reviewed the present situation, her knowledge about sensory issues and their impact on arousal regulation and mealtime participation may lead her to refer to the OT. The opposite referral could also occur; for example, the OT may identify parental anxiety and inadequate family dynamics around mealtimes as one of the underlying issues to the child's eating problem and refer the family to a psychologist.

### Gastrointestinal processes

Literature suggests a higher prevalence of gastrointestinal issues in children with ASD, with constipation and gastroesophageal reflux being among the most frequent (Ibrahim *et al.*, 2009). Gastroesophageal reflux and constipation should be considered in the assessment, as both are reported to coexist with eating problems (Ibrahim *et al.*, 2009). In clinical practice, we often observe that children who experience discomfort related to the feeding process develop a negative relationship with food and show little motivation to eat (Kerzner *et al.*, 2015). The intervention should include the treatment of the digestive disorder and consideration of the refusal to eat. The expectation that everything will be fine once the medical problem has been resolved is rarely met (Zangen *et al.*, 2003), and the collaboration between the gastroenterologist and the rest of the team should begin early on.

### Nutrition

Reports of the nutritional status of children with ASD indicate comparable intakes of energy, carbohydrates and fats when compared to typically developing peers (Sharp *et al.*, 2013). However, closer examination indicates deficits in calcium and protein intake and a higher number of nutritional deficits among children with ASD (Sharp *et al.*, 2018). Relying exclusively on anthropometric parameters such as weight, height and body mass index to assess health status is not sufficient. The Three-Day Food Diary (Cornish, 2002) is a tool that allows professionals in the field of nutrition to measure the nutritional consumption of the child and compare it to a reference value. This type of tool is used to measure the quantity and variety of foods consumed. However, the information gathered by the means of a food diary gives a limited vision of the eating problem (Nadon *et al.*, 2008). If food variety and/or quantity is found to be limited, further interventions with nutritionists and therapists can be implemented to expand food variety and improve nutritional status.

The analysis of the nutritional needs of the child, together with the assessment of oral sensorimotor skills, allows the team to set a diet that takes into account the skills of the child, the nutritional value, texture, taste and presentation of food (Beaudry-Bellefeuille *et al.*, 2015). For example, the nutritionist may prioritise fibre and could recommend foods such as broccoli and strawberries. However, the OT may consider the sensory properties of these foods to be too difficult for the child and discuss with the nutritionist the need to identify high-fibre foods with a more homogenous texture. Children with ASD may also have to follow a specific diet that their family chooses because of ethical or religious reasons (e.g. dairy-free or meat-free diet). In these cases, the contribution of

a nutritionist is equally important. Through collaboration, we aim to provide consensual recommendations that are in accordance with the family's diet, the child's nutritional needs, sensorimotor abilities and general preferences to avoid recommending foods that will likely be refused and cause more mealtime problems.

### Oro-motor abilities

Oro-motor skills for eating are a complex set of fine motor skills, which are mostly established in typical development by three years (Morris and Klein, 2000). Their progression is embedded in the context of both the child's gross motor and sensory development, and difficulties in this area can lead to food refusal and selectivity (Morris and Klein, 2000). The literature outlines oro-motor difficulties in children with ASD (Nadon *et al.*, 2013) and in children demonstrating oral aversions, avoidance or fear of eating (Godoy *et al.*, 2019). Difficulty progressing to challenging food textures, gagging, food loss, poor mouth clearance, swallowing issues or drooling may all be manifestations of poor oro-motor control (Smile *et al.*, 2020; Sharp *et al.*, 2013). In case such difficulties are noticed, assessment of the child's oral motor skills is vital.

This area has been extensively researched and developed clinically within the fields of speech therapy and occupational therapy (Marcus and Breton, 2013; Morris and Klein, 2000). Our evaluation is based on extensive knowledge of oral motor development (Morris and Klein, 2000) and the observation of a meal that involves preferred and non-preferred foods. Therapists who are knowledgeable in the area of oro-motor issues can build an intervention plan to improve the oral skills that may be at the root of poor control of food, ineffective bolus formation and inefficient chewing. Clinically, we often observe that children with oro-motor issues will refuse foods which require refined oral skills and develop strong preferences for the foods that are less challenging motorically. Making a list of preferred and non-preferred foods is a way to identify a possible pattern linked to motor skill. For example, if all preferred foods are mashed, soft and/or dissolvable and non-preferred foods require refined chewing skills, this may be an indication of oro-motor issues. In these cases, safety is always a concern and foods must be carefully chosen to avoid aspiration. Dissolvable solids such as crackers are often a good option. Feeders that hold the food in a gauze-like pouch can also be used, allowing the child to practice chewing skills while safely securing the food inside the feeder.

### Sensory functions

A meal is a complex sensory experience consisting of the foods with their appearance, odours, textures and tastes, as well as the presence of others. When considering sensory functions, several aspects must be assessed. Sensory reactivity, sensory perception, praxis, postural control and bilateral integration have all been identified to be part of sensory functions (Ayes, 2004). Issues in sensory functions are common in children with ASD and may potentially impact feeding (Zobel-Lachiusa *et al.*, 2015).

Sensory hyper-reactivity has been widely identified to be among one of the main factors related to food rejection and food selectivity in children with ASD (Zobel-Lachiusa *et al.*, 2015). Sensory hyper-reactivity is also common in children

with a history of gastroesophageal reflux (Davis *et al.*, 2013), a common GI issue in children with ASD (Ibrahim *et al.*, 2009). Making a list of preferred and non-preferred foods can also be extremely useful to identify sensory reactivity issues. For example, if preferred foods are homogenous, dry and/or smooth in texture and non-preferred foods are of mixed, viscous or lumpy textures, this could be an indication of issues in sensory reactivity. The use of standardised questionnaire such as the sensory processing measure (Parham *et al.*, 2007) is a key component of the assessment of sensory reactivity issues.

Sensory perception must also be considered. Bennetto *et al.* (2007) found that children with suggesting ASD may struggle to identify taste and olfactory sensations suggesting that issues in sensory perception contribute to eating problems among this population. Somatosensory perception difficulties can also potentially impact eating. Multiple studies have reported a relationship between somatosensory discrimination and praxis (Ayres, 2004). Research has shown that many individuals with ASD have praxis and imitation difficulties, including orofacial imitation (Mostofsky *et al.*, 2006). Furthermore, somatosensory perception deficits, in combination with issues in praxis, are reported to be frequent in children with ASD (Roley *et al.*, 2015). A common observation related to this type of issue is the lack of ability to localise food in the mouth and organise tongue movements to handle the food. Children may prefer food that is soft and homogeneous (e.g. mashed foods), not because of motor problems as such but because of difficulty in locating the food in the mouth and planning oral movements accordingly. Clinically, we observe that children benefit from intense oral sensorimotor activities such as biting on vibrating or textured oral toys, becoming more aware of their oral cavity and better equipped to handle a variety of textures. Choosing foods that can be safely handled with limited intraoral perception and praxis is of utmost importance.

Children, from an early age, are expected to adapt to the family's routine and learn the "mealtime rules" by modelling the eating behaviours of parents and siblings (Birch *et al.*, 1989). Research has shown that modelling healthy eating habits can have a positive effect on expanding children's dietary preferences (Birch *et al.*, 1989). Therefore, difficulty in imitating other people's behaviour could compromise the broadening of the eating repertoire (Nadon *et al.*, 2011), and assessment of somatosensory perception and praxis is warranted in children who fail to imitate parents and peers when trying new foods. Assessment tools such as the Sensory Integration and Praxis Tests (Ayres, 2004) are useful to identify underlying sensory perception and praxis issues that may be impacting mealtime participation.

Efficient processing of vestibular and proprioceptive input is necessary for general motor skills such as trunk control for sitting upright, a key component of mealtime participation, and must therefore be considered as part of the eating assessment (Marcus and Breton, 2013). Deficits in postural stability and motor coordination in individuals with ASD are well-documented (Flanagan *et al.*, 2012) and should be assessed when sitting at the table and using hands for self-feeding are problematic. In children with ASD, difficulties with vestibular-proprioceptive processing are often manifested as moving excessively and can easily be misinterpreted as a behavioural issue. The OT intervention to improve mealtime participation

will often include adaptation of seating options and direct therapy for the underlying vestibular-proprioceptive issues.

OTs have developed expertise in sensorimotor deficits that impact participation in activities of daily living, and children who show issues in this area may benefit from an in-depth assessment of their sensory functions.

### Respiratory processes

Issues with breathing are reported to be present in up to 25% of children with autism (Williams *et al.*, 2004). The accumulation of secretions, breathing by the mouth and respiratory effort can impact the feeding process (Trabalon and Schaal, 2012). In these cases, the child is referred to a medical specialist (allergist, otorhinolaryngologist) for assessment and treatment.

### Early feeding behaviour

The findings from the Avon Longitudinal Study of Parents and Children (Emond *et al.*, 2010) indicate that children with a subsequent ASD diagnosis were more commonly described as "slow feeders" by parents at six months and had a slow transition to solid foods. Also, at 15 to 54 months, it was noted that toddlers with ASD were "difficult to feed" and "very choosy" eaters (Emond *et al.*, 2010). In another study, Brisson *et al.* (2012) collected family videos of 48 children with ASD and 46 typically developing children and studied retrospectively how often the babies opened their mouth in anticipation of the feeding spoon. Researchers observed that typically developing infants who had initial anticipation difficulties quickly learned to successfully anticipate. However, this did not happen with infants who later received an ASD diagnosis.

From a clinical perspective, there is literature suggesting that an early assessment followed by consistent monitoring of infant eating behaviour is crucial. Although exploring some of the early biological and behavioural markers of ASD can be invasive (e.g. brain imaging and eye-tracking techniques), assessing the eating behaviour of children from infancy is a good practice that will promote the overall health and the development of children, but if eating problems are accompanied by any other early symptoms related to ASD, it can raise clinicians and carers attention and perhaps lead to seeking an earlier diagnosis or intervention (van't Hof, *et al.*, 2020).

### Communication and social skills

Children with ASD are more likely to present with developmental delays in the areas of speech and social interaction (APA, 2013). Mealtime is one of the key social interaction moments both for the family and the child. Consequently, the existence of social and communication difficulties may make mealtime a stressful time and may compromise the ability to effectively communicate needs around food (Williams *et al.*, 2000). In these cases, therapists or any other teaching staff working with the child should prioritise the expansion of the child's mealtime vocabulary so that they can better communicate their food preferences, or any sensory or intestinal discomfort certain foods may cause them.

### Psychological factors affecting eating in children with autism spectrum disorders

Children with ASD commonly engage in repetitive and ritualistic behaviour (Boyd *et al.*, 2010). Consequently, they are more likely than typically developing children to insist on a ritualistic mealtime, such as eating the same food every day (Schreck *et al.*, 2004). Children with ASD may also demonstrate insistence for sameness, which can result in a preference to use certain utensils or follow certain routines during mealtime. Another factor that can compromise eating is anxiety, a frequent co-occurring diagnosis in this population (MacNeil *et al.*, 2009), which can decrease appetite (Bryant-Waugh *et al.*, 2010). Lack of appetite may mistakenly be perceived as extreme food selectivity or “fussiness” in children with ASD.

Psychologists should be aware that repetitive and ritualistic behaviour may also be demonstrated in the eating behaviour of children with ASD. In these cases, the psychologist needs to work with the child with ASD to specifically decrease their anxiety around mealtime and/or develop trust towards foods that they are less willing to try and assisting them in the introduction of these foods into their diet (Dial *et al.*, 2020).

### Summary and conclusions

Children with ASD and eating problems have complex combinations of medical, nutritional, feeding skills and/or psychosocial issues (Smile *et al.*, 2020). Commonly used terms such as “picky eater”, “food refusal” and “food selectivity” seem to imply that these are voluntary behaviours when they are very likely to appear because of unidentified medical, psychological and/or sensorimotor problems that make the process of eating challenging or painful (Williams *et al.*, 2010).

Reports of interventions that include comprehensive assessment, analysis of objective data and clinical reasoning to identify underlying issues that compromise participation in daily activities are showing positive outcomes in children with ASD (Schaaf *et al.*, 2014). Similarly, this approach to assessment and intervention has also shown promising results in children with eating problems (Beaudry-Bellefeuille *et al.*, 2015; Gándara-Gafo *et al.*, 2021). Focussing on the multiple underlying mechanisms that may impact eating can help us better understand the eating problems of this population. Once the underlying factors are identified, the professionals of the multidisciplinary team who are best equipped to tackle the challenges can design the intervention in collaboration with the family (Goday *et al.*, 2019). Pleasurable participation in mealtimes for the child and family, as well as the development of healthy eating habits that support growth and development, are the goals of the intervention. OTs, speech therapists, nutritionists, psychologists and several medical specialties may all be necessary to carry out a personalised assessment and intervention when dealing with children with ASD who face eating problems.

### References

American Psychiatric Association (2013), “Diagnostic and statistical manual of mental disorders (DSM-5®)”, American Psychiatric Pub.

- Ayres, A.J. (2004), *Sensory Integration and Praxis Tests Manual*, updated edition. Western Psychological Services, Los Angeles, CA.
- Bandini, L.G., Anderson, S.E., Curtin, C., Cermak, S., Evans, E.W., Scampini, R. and Must, A. (2010), “Food selectivity in children with autism spectrum disorders and typically developing children”, *The Journal of Pediatrics*, Vol. 157 No. 2, pp. 259-264.
- Beaudry-Bellefeuille, I., Ramos-Polo, E., Suarez-González, M., Moriyon-Iglesias, T. and Gandara-Gafo, B. (2015), “Colaboración multidisciplinaria en el tratamiento de un niño de 5 años de edad con desnutrición”, *Acta Pediátrica Española*, Vol. 73 No. 5, pp. 127-133.
- Bennetto, L., Kuschner, E.S. and Hyman, S.L. (2007), “Olfaction and taste processing in autism”, *Biological Psychiatry*, Vol. 62 No. 9, pp. 1015-1021, doi: [10.1016/j.biopsych.2007.04.019](https://doi.org/10.1016/j.biopsych.2007.04.019).
- Birch, L.L., McPhee, L., Sullivan, S. and Johnson, S. (1989), “Conditioned meal initiation in young children”, *Appetite*, Vol. 13 No. 2, pp. 105-113, doi: [10.1016/0195-6663\(89\)90108-6](https://doi.org/10.1016/0195-6663(89)90108-6).
- Boyd, B.A., Baranek, G.T., Sideris, J., Poe, M.D., Watson, L. R., Patten, E. and Miller, H. (2010), “Sensory features and repetitive behaviors in children with autism and developmental delays”, *Autism Research : Official Journal of the International Society for Autism Research*, Vol. 3 No. 2, pp. 78-87.
- Brisson, J., Warreyn, P., Serres, J., Foussier, S. and Adrien-Louis, J. (2012), “Motor anticipation failure in infants with autism: a retrospective analysis of feeding situations”, *Autism*, Vol. 16 No. 4, pp. 420-429.
- Broder-Fingert, S., Brazauskas, K., Lindgren, K., Iannuzzi, D. and Van Cleave, J. (2014), “Prevalence of overweight and obesity in a large clinical sample of children with autism”, *Academic Pediatrics*, Vol. 14 No. 4, pp. 408-414.
- Bryant-Waugh, R., Markham, L., Kreipe, R.E. and Walsh, B.T. (2010), “Feeding and eating disorders in childhood”, *International Journal of Eating Disorders*, Vol. 43 No. 2, p. 9, doi: [10.1002/eat.20795](https://doi.org/10.1002/eat.20795).
- Cornish, E. (2002), “Gluten and casein free diets in autism: a study of the effects on food choice and nutrition”, *Journal of Human Nutrition and Dietetics*, Vol. 15 No. 4, pp. 261-269.
- Davis, A.M., Bruce, A.S., Khasawneh, R., Schulz, T., Fox, C. and Dunn, W. (2013), “Sensory processing issues in young children presenting to an outpatient feeding clinic: a retrospective chart review”, *Journal of Pediatric Gastroenterology & Nutrition*, Vol. 56 No. 2, p. 156.
- Dial, L.A., Emley, E., Koerten, H.R., Waite, T.C. and Musher-Eizenman, D.R. (2020), “A “mindfulness intervention for food neophobia among preschoolers”, *Early Childhood Education Journal*, Vol. 48 No. 1, pp. 117-126.
- Emond, A., Emmett, P., Steer, C. and Golding, J. (2010), “Feeding symptoms, dietary patterns, and growth in young children with autism spectrum disorders”, *PEDIATRICS*, Vol. 126 No. 2, pp. e337-e342.
- Flanagan, J.E., Landa, R., Bhat, A. and Bauman, M. (2012), “Head lag in infants at risk for autism: a preliminary study”, *American Journal of Occupational Therapy*, Vol. 66 No. 5, pp. 577-585, doi: [10.5014/ajot.2012.004192](https://doi.org/10.5014/ajot.2012.004192).

- Florence, M.D., Asbridge, M. and Veugelers, P.J. (2008), "Diet quality and academic performance", *Journal of School Health*, Vol. 78 No. 4, pp. 209-215.
- Gándara-Gafo, B., Moriyón-Iglesias, T. and Beaudry-Bellefeuille, I. (2021), "Occupational therapy assessment and intervention of a 22-month-old girl with feeding refusal", *Journal of Behavioral Health*, Vol. 10 No. 1, pp. 1-3.
- Goday, P.S., Huh, S.Y., Silverman, A., Lukens, C.T., Dodrill, P., Cohen, S.S. and Phalen, J.A. (2019), "Pediatric feeding disorder: consensus definition and conceptual framework", *Journal of Pediatric Gastroenterology & Nutrition*, Vol. 68 No. 1, pp. 124-129, doi: [10.1097/MPG.0000000000002188](https://doi.org/10.1097/MPG.0000000000002188).
- Ibrahim, S.H., Voigt, R.G., Katusic, S.K., Weaver, A.L. and Barbaresi, W.J. (2009), "Incidence of gastrointestinal symptoms in children with autism: a population-based study", *PEDIATRICS*, Vol. 124 No. 2, pp. 680-686.
- Kanner, L. (1943), "Autistic disturbances of affective contact", *Nervous Child*, Vol. 2 No. 3, pp. 217-250.
- Kerzner, B., Milano, K., MacLean, W.C., Berall, G., Stuart, S. and Chatoor, I. (2015), "A practical approach to classifying and managing feeding difficulties", *PEDIATRICS*, Vol. 135 No. 2, pp. 344-353.
- Kodak, T. and Piazza, C.C. (2008), "Assessment and behavioral treatment of feeding and sleeping disorders in children with autism spectrum disorders", *Child and Adolescent Psychiatric Clinics of North America*, Vol. 17 No. 4, pp. 887-905.
- Kuschner, E.S., Eisenberg, I.W., Orionzi, B., Simmons, W.K., Kenworthy, L., Martin, A. and Wallace, G.L. (2015), "A preliminary study of self-reported food selectivity in adolescents and young adults with autism spectrum disorder", *Research in Autism Spectrum Disorders*, Vol. 15, pp. 53-59.
- MacNeil, B.M., Lopes, V.A. and Minnes, P.M. (2009), "Anxiety in children and adolescents with autism spectrum disorders", *Research in Autism Spectrum Disorders*, Vol. 3 No. 1, pp. 1-21.
- Marcus, S. and Breton, S. (2013), *Infant and Child Feeding and Swallowing: occupational Therapy Assessment and Intervention*, American Occupational Therapy Association Press, North Bethesda, MD.
- Mari-Bauset, S., Zazpe, I., Mari-Sanchis, A., Llopis-González, A. and Morales-Suárez-Varela, M. (2014), "Food selectivity in autism spectrum disorders: a systematic review", *Journal of Child Neurology*, Vol. 29 No. 11, pp. 1554-1561.
- Mari-Bauset, S., Llopis-González, A., Zazpe-García, I., Mari-Sanchis, A. and Morales-Suárez-Varela, M. (2015), "Nutritional status of children with autism spectrum disorders (ASDs): a case-control study", *Journal of Autism and Developmental Disorders*, Vol. 45 No. 1, pp. 203-212.
- Morris, S.E. and Klein, M.D. (2000), *Pre-Feeding Skills: A Comprehensive Resource for Mealtime Development*, Pro-ed., Austin.
- Mostofsky, S.H., Dubey, P., Jerath, V.K., Jansiewicz, E.M., Goldberg, M.C. and Denckla, M.B. (2006), "Developmental dyspraxia is not limited to imitation in children with autism spectrum disorders", *Journal of the International Neuropsychological Society*, Vol. 12 No. 03, pp. 314-326.
- Nadon, G., Feldman, D. and Gisel, E. (2013), "Feeding issues associated with the autism spectrum disorders", in Fitzgerald, M. (Ed.), *Recent Advances in Autism Spectrum Disorders-Volume I*, IntechOpen, doi: [10.5772/53644](https://doi.org/10.5772/53644).
- Nadon, G., Ehrmann-Feldman, D. and Gisel, E. (2008), "Revue des méthodes utilisées pour évaluer l'alimentation des enfants présentant un trouble envahissant du développement", *Archives de Pédiatrie*, Vol. 15 No. 8, pp. 1332-1348, doi: [10.1016/j.arcped.2008.04.022](https://doi.org/10.1016/j.arcped.2008.04.022).
- Nadon, G., Feldman, D.E., Dunn, W. and Gisel, E. (2011), "Association of sensory processing and eating problems in children with autism spectrum disorders", *Autism Research and Treatment*, Vol. 2011, doi: [10.1155/2011/541926](https://doi.org/10.1155/2011/541926).
- Parham, L., Ecker, C., Miller-Kuhaneck, H., Henry, D. and Glennon, T. (2007), *Sensory Processing Measure*, Western Psychological Services, Los Angeles.
- Roley, S.S., Mailloux, Z., Parham, L.D., Schaaf, R.C., Lane, C.J. and Cermak, S. (2015), "Sensory integration and praxis patterns in children with autism", *American Journal of Occupational Therapy*, Vol. 69 No. 1, pp. 6901220010p1-6901220010p8.
- Schaaf, R.C., Benevides, T., Mailloux, Z., Faller, P., Hunt, J., van Hooydonk, E. and Kelly, D. (2014), "An intervention for sensory difficulties in children with autism: a randomized trial", *Journal of Autism and Developmental Disorders*, Vol. 44 No. 6, pp. 1493-1506.
- Schreck, K.A., Williams, K. and Smith, A.F. (2004), "A comparison of eating behaviors between children with and without autism", *Journal of Autism and Developmental Disorders*, Vol. 34 No. 4, pp. 433-438.
- Sharp, W.G., Berry, R.C., McCracken, C., Nuhu, N.N., Marvel, E., Saulnier, C.A. and Jaquess, D.L. (2013), "Feeding problems and nutrient intake in children with autism spectrum disorders: a meta-analysis and comprehensive review of the literature", *Journal of Autism and Developmental Disorders*, Vol. 43 No. 9, pp. 2159-2173.
- Sharp, W.G., Postorino, V., McCracken, C.E., Berry, R.C., Criado, K.K., Burrell, T.L. and Scahill, L. (2018), "Dietary intake, nutrient status, and growth parameters in children with autism spectrum disorder and severe food selectivity: an electronic medical record review", *Journal of the Academy of Nutrition and Dietetics*, Vol. 118 No. 10, pp. 1943-1950, doi: [10.1016/j.jand.2018.05.005](https://doi.org/10.1016/j.jand.2018.05.005).
- Smile, S., Raffaele, C. and Perlin, R. (2020), "Re-imagining the physicians' role in the assessment of feeding challenges in children with autism spectrum disorder", *Paediatrics and Child Health*, doi: [10.1093/pch/pxaa008](https://doi.org/10.1093/pch/pxaa008).
- Trabalon, M. and Schaal, B. (2012), "It takes a mouth to eat and a nose to breathe: abnormal oral respiration affects neonates' oral competence and systemic adaptation", *International Journal of Pediatrics*, Vol. 2012, doi: [10.1155/2012/207605](https://doi.org/10.1155/2012/207605).
- Twachtman-Reilly, J., Amaral, S.C. and Zebrowski, P.P. (2008), "Addressing feeding disorders in children on the autism spectrum in school-based settings: physiological and behavioral issues", *Language, Speech, and Hearing Services in Schools*, Vol. 39 No. 2, pp. 261-272, doi: [10.1044/0161-1461\(2008/025\)](https://doi.org/10.1044/0161-1461(2008/025)).
- van't Hof, M., Ester, W.A., van Berckelaer-Onnes, I., Hillegers, M.H., Hoek, H.W. and Jansen, P.W. (2020),

- “Do early-life eating habits predict later autistic traits? Results from a population-based study”, *Appetite*, Vol. 156, p. 104976.
- Williams, P.G., Dalrymple, N. and Neal, J. (2000), “Eating habits of children with autism”, *Pediatric Nursing*, Vol. 26 No. 3, pp. 259-265.
- Williams, K.E., Field, D.G. and Seiverling, L. (2010), “Food refusal in children: a review of the literature”, *Research in Developmental Disabilities*, Vol. 31 No. 3, pp. 625-633.
- Williams, G., Sears, L. and Allard, A. (2004), “Sleep problems in children with autism”, *Journal of Sleep Research*, Vol. 13 No. 3, pp. 265-268.
- Zangen, T., Ciarla, C., Zangen, S., Di Lorenzo, C., Flores, A.F., Cocjin, J. and Hyman, P.E. (2003),

- “Gastrointestinal motility and sensory abnormalities may contribute to food refusal in medically fragile toddlers”, *Journal of Pediatric Gastroenterology and Nutrition*, Vol. 37 No. 3, pp. 287-293.
- Zobel-Lachiusa, J., Andrianopoulos, M.V., Mailloux, Z. and Cermak, S.A. (2015), “Sensory differences and mealtime behaviour in children with autism”, *American Journal of Occupational Therapy*, Vol. 69 No. 5, pp. 6905185050p1-6905185050p8, doi: [10.5014/ajot.2015.016790S](https://doi.org/10.5014/ajot.2015.016790S).

### Corresponding author

Isabelle Beaudry-Bellefeuille can be contacted at: [ibbergo@gmail.com](mailto:ibbergo@gmail.com)